

# Surface water monitoring at 30 m resolution; prototyping new Copernicus Global Land Service products

J.-F. Pekel, A. Cottam, M. Clerici, A. Belward, G. Dubois, E. Bartholome (EC JRC) and N.Gorelick (GEE)



Landsat 8
graphic courtesy USGS



Sentinel 2 graphic courtesy ESA









# **Global Land Service**

Q search this site

Home Products FAQ Helpdesk Links News & events About the global land service

Copernicus Land service


#### Home

The Global Land Component in the framework of GMES Initial Operations (GIO) is earmarked as a component of the Land service to operate "a multi-purpose service component" that will provide a series bio-geophysical products on the status and evolution of land surface at global scale. Production and delivery of the parameters are to take place in a timely manner and are complemented by the constitution of long term time series.

Read more

## **Special images**

Space and Time Syntheses (April - September 2013, combine start/duration/intensity/ extension of FAPAR values) produced using GIO data to present draught in Slovenia.



### Go to product:

√ - Choose Product -

Burnt Area (BA)

Dry Matter Productivity (DMP)

Fraction of Absorbed Photosynthetically Active Radiation (FAPAR)

Fraction of green Vegetation Cover (FCover)

Leaf Area Index (LAI)

Normalized Difference Vegetation Index (NDVI)

Vegetation Condition Index (VCI)

Vegetation Productivity Index (VPI)

Land Surface Temperature (LST)

Surface Albedo (SA)

Top Of Canopy Reflectances (TOC-r)

Water Bodies (WB)


World Water Day 2014 and the importance of Water Bodies products!

#### News published:

1 month 1 week ago



- 1. Human wellbeing and economy water supply
- 2. Human security flood and drought monitoring
- 3. Essential Climate Variables lake area, river discharge (modeling, impacts, adaptation)
- **4. Food security** irrigation, rangeland management
- **5. Human and animal health** pollution, poisons, disease and vectors
- **6. Biodiversity** wetland, lacustrine and riverine ecosystems, invasive species and migratory corridors
- 7. Coastline, lake, river boundary changes infrastructure planning, insurance, reference databases





- 1. Permanent water surfaces
- 2. Seasonal water surfaces (and dates)
- 3. Ephemeral water surfaces (and dates)
- 4. New water surfaces
- 5. Never water surfaces



Katse Dam, Lesotho 2005



**Badwater Death Valley, USA 2005** 



Lake Maggiore, Italy 2009




Okavango delta, Botswana 2006












Corvallis Landsat 6<sup>th</sup> July 2014 Courtesy USGS / NASA





# Surface water detection method

- 1. Based on Hue/Saturation/Value colour model rather than RGB
- 2. Sensor neutral used on L5, L7, L8 and MODIS
- 3. Uses Google Earth Engine for processing

4. Uses tools built on top of GEE - for spectral library development,

classification and validation

5. Uses a number of masks:

Cloud


**Snow** 

Vegetation

Slope


Bare rock

Temperature masks





# **Surface water examples**




Landsat imagery (courtesy USGS / NASA) extracted from Google Earth Engine





## Surface water detected




Landsat imagery (courtesy USGS / NASA) extracted from Google Earth Engine





Source; Andrew Cottam, J-F Pekel (JRC) Noel Gorelick (GEE)



# **Inter Annual variation**












# **Intra Annual variation**







- 1. Global land surface process monitoring at Landsat-class resolution is a reality, but we are working at the limits of the system...
- high quality radiometry and calibration are needed; which also allows multi-platform/sensor analysis...
- 3. which leads to better inter and intra annual temporal sampling...
- and thus completely new products and services that meet the needs of many end-users



Grand Renaissance Dam Blue Nile, Ethiopia 15 October 2000



Grand Renaissance Dam Blue Nile, Ethiopia 4 October 2011



Grand Renaissance Dam Blue Nile, Ethiopia 19 October 2013

